



Deworming Protocol



Fecal egg counts (FEC) provide us with valuable information that help us evaluate the efficacy of dewormers being used and determine the best deworming treatment plan for your horse. Spring and Fall are peak transmission time, by doing a FEC during these times and following a deworming protocol we can minimize pasture contamination and prevent parasitic disease and resistance.

Adult LOW Shedder Treatment Plan

SPRING	FALL
RUN A FECAL TEST	RUN A FECAL TEST
DEWORMER TO USE: <ul style="list-style-type: none"> Moxidectin (QUEST) Ivermectin for miniature horses 	DEWORMER TO USE: <ul style="list-style-type: none"> Ivermectin with Praziquantel (EQUIMAX)

Adult HIGH Shedder Treatment Plan

MONTH	FECAL TEST?	DEWORMER TO USE
January	NO	<ul style="list-style-type: none"> Ivermectin
March	YES	<ul style="list-style-type: none"> Moxidectin (QUEST) Ivermectin for miniature horses
May	NO	<ul style="list-style-type: none"> Double dose of Pyrantel (STRONGID)
July	NO	<ul style="list-style-type: none"> Ivermectin with Praziquantel (EQUIMAX)
October	YES	<ul style="list-style-type: none"> Moxidectin (QUEST) Ivermectin for miniature horses



Dosage is based on weight!

Foal Treatment Plan

Age	Treatment Plan	Brand Name of Dewormer
2 Months	Fenbendazole (10mg/kg)	Panacur
3 Months	Pyrantel	Strongid
4 Months	Ivermectin	Zimecterin, Bimectin, Ivermectin Paste
4 ½ Months	Fecal Test!	
6 Months	Pyrantel	Strongid
8 Months	Ivermectin with Praziquantel	Equimax, Zimecterin Gold
10 Months	Pyrantel	Strongid
12 Months	Moxidectin	Quest for 700 lbs and up & Panacur Power Pak for under 700 lbs